

Referenser

1. Polatin et al. Psychiatric illness and cronic low back pain. The mine and the spine-witch goes first?. Spine 18;66-71, 1993.
2. Axis I and Axis II psychiatric disorders in patients with fibromyalgia , Faruk Uguz M.D.a, Erdinç Çiçek M.D.a, Ali Salli M.D.b, Ali Yavuz Karahan M.D.b, İlknur Albayrak M.D.b, Nazmiye Kaya M.D.a and Hatice Uğurlu M.D.b General Hospital Psychiatry, vol 32,January-February 2010, Pages 105-107
3. Bair MJ, et al. *Arch Intern Med.* 2003;**163**:2433–2445.
4. Turk och Okifuji, Assessment of patients' reporting of pain: an integrated perspective, *Lancet* 1999, vol 353, sid 1784-88
5. *Psychosom Med* 2008
6. Von Knorring, L Ekselius: >Idiopathic pain and depression. Quality of Life Res 1994;3:S57-S68.
7. Simon GE, et al. *N Engl J Med.*1990;341:1329-1335
8. Wittchen et al *J Clin Psychiatry.* 2002;63(suppl8):24-34. Brandenburg et al 2005.
9. Allgulander, Nilsson Rikstäckande primärvårdsstudie:Var fjärde patientlider av ångest och depressionLäkartidningent 2003 Bodlund O. Anxiety and depression as a hidden problem in primary health care. Only one case in four identified [summary]. *Läkartidningen* 1997; 94: 4612-4, 4617-8.
10. Lundberg M. Kinesiophobia - various aspects of moving with musculoskeletal pain.