

Referenser

1. Socialstyrelsen. Diagnostik och läkemedelsbehandling vid adhd. Förekomst, trend och könsskillnader. Stockholm: Socialstyrelsen, 2023.
2. Adler M, Bengtsson J, Rück C. Låt inte ovaliderade diagnoser få fäste innan evidens finns. Läkartidningen 2024; 9-10.
3. Widiger TA. A dimensional model of psychopathology. Psychopathology 2005; 38(4): 211-4.
4. Das D, Cherbuin N, Butterworth P, Anstey KJ, Easteal S. A population-based study of attention deficit/hyperactivity disorder symptoms and associated impairment in middle-aged adults. PloS one 2012; 7(2): e31500.
5. Larsson H, Anckarsater H, Råstam M, Chang Z, Lichtenstein P. Childhood attention-deficit hyperactivity disorder as an extreme of a continuous trait: a quantitative genetic study of 8,500 twin pairs. Journal of child psychology and psychiatry, and allied disciplines 2012; 53(1): 73-80.
6. Heidbreder R. ADHD symptomatology is best conceptualized as a spectrum: a dimensional versus unitary approach to diagnosis. Attention deficit and hyperactivity disorders 2015; 7(4): 249-69.
7. Arildskov TW, Thomsen PH, Sonuga-Barke EJS, Lambek R, Østergaard SD, Virring A. Is Attention-Deficit/Hyperactivity Disorder (ADHD) a Dimension or a Category? What Does the Relationship Between ADHD Traits and Psychosocial Quality of Life Tell Us? Journal of Attention Disorders 2024; 28(7): 1035-44.